

For remove and collect of Leukocytes



You can remove Leukocytes only by filtering, NO centrifuge is necessary. Moreover, you can collect trapped Leukocytes in the filter!!




You can trap, remove and collect leukocytes specifically from whole blood by Leuko filter.

Cat No.	ltem	Packaging
1883-110LC	LeukoCatch [®] II, 10 mL, Column Type	Sterilized 5 pieces / bag

The leukocyte count after filtration by LeukoCatch II

6 out of 20 fields randomly obtained from the microscopic abservation of a mouse blood



*Data provider: Mie University, Faculty of Medicine, The Laboratory of Medical Zoology



①Set LeukoCatch II into a 50 mL centrifuge tube, and apply 10 mL of blood. (5 mL whole blood + 5 mL PBS(-))



②Filtering starts with natural dropping force. Filtering time is approx. 10 min.



③Leuko Filter traps leykocytes. Consequently, leukocytes are almost removed from the filtrate.

Collect Leukocytes which are trapped

by filter of product.

There are concave side and convex side in Leuko Filter

Convex side is upper side

After filtration of sample, a lot of leukocytes are attached to convex side (upper side).




Concave side is back side Leuko Filter is dented in center of the back side



4) Add 2 mL of PBS(-) in order to wash the Leuko Filter trapped leukocytes. Repeat 1 time.



⑤ Pick up Leuko Filter with a pair of tweezers and remove Leuko Filter.




Pay attention NOT to reverse the face of Leuko Filter.



@Pool the 5 mL of culture medium into a Φ60 mm dish.

Put Leuko Filter in the culture medium.



⑦Put the dish in the CO2 incubator. (37 degree Celsius)



® After 24 houres, it can collect about 60% leukocytes from whole blood without centrifugation.

Note: As cited from an article in Advances in Bioscience and Biotechnology, 2012.

"Nevertheless, these leukocyte separation procedures require laboratory-based professional and time-consuming procedures using instrumentation that is not routinely available at the bedside or in a common clinical facility. Moreover, their protocols are too complex to apply to an automated machine to treat hundreds of samples simultaneously."



Manufactured and sold by:

FUKAEKASEI CO., LTD.

Head Office: 2-2-7 Murotani, Nishi-ku, Kobe 651-2241, JAPAN TEL +81-78-991-4477

FAX +81-78-991-4491 E-mail: info@watson.co.jp

https://watsonbiolab.com